

Provider: _____ Clinic: _____



My Asthma Action Plan

Name: _____ DOB: ____ / ____ / ____

Severity Classification: Intermittent Mild Persistent Moderate Persistent Severe Persistent

Asthma Triggers (list): _____

Peak Flow Meter Personal Best: _____

Green Zone: Doing Well

Symptoms: Breathing is good – No cough or wheeze – Can work and play – Sleeps well at night

Peak Flow Meter _____ (more than 80% of personal best)

Flu Vaccine—Date received: _____ Next flu vaccine due: _____ COVID19 vaccine—Date received: _____

Control Medicine(s)	Medicine	How much to take	When and how often to take it
_____	_____	_____	_____
_____	_____	_____	_____

Physical Activity

- Use Albuterol/Levalbuterol _____ puffs, 15 minutes before activity
 with all activity when you feel you need it

Yellow Zone: Caution

Symptoms: Some problems breathing – Cough, wheeze, or tight chest – Problems working or playing – Wake at night

Peak Flow Meter _____ to _____ (between 50% and 79% of personal best)

Quick-relief Medicine(s) Albuterol/Levalbuterol _____ puffs, every 20 minutes for up to 4 hours as needed

Control Medicine(s) Continue Green Zone medicines

Add _____ Change to _____

You should feel better within 20-60 minutes of the quick-relief treatment. If you are getting worse or are in the Yellow Zone for more than 24 hours, THEN follow the instructions in the RED ZONE and call the doctor right away!

Red Zone: Get Help Now!

Symptoms: Lots of problems breathing – Cannot work or play – Getting worse instead of better – Medicine is not helping

Peak Flow Meter _____ (less than 50% of personal best)

Take Quick-relief Medicine NOW! Albuterol/Levalbuterol _____ puffs, _____ (how frequently)

Call 911 immediately if the following danger signs are present:

- Trouble walking/talking due to shortness of breath
- Lips or fingernails are blue
- Still in the Red Zone after 15 minutes

Emergency Contact Name _____ Phone (_____) _____ - _____

Date: ____ / ____ / ____

Plan de acción de asma

Nombre: _____ Fecha de nacimiento: ____ / ____ / ____

Clasificación de severidad: Intermitente Persistente leve Persistente moderado Persistente severa

Desencadenantes del asma (lista): _____

Mejor marca personal del medidor de flujo: _____

Zona verde: Anda bien

Síntomas: La respiración es buena – Sin tos, ni sibilancias – Puede trabajar y jugar – Duerme bien en la noche

Medidor de flujo _____ (más del 80% de la mejor marca personal)

Medicamento/s de control	Medicamento	Cuánto tomar	Cuándo y con qué frecuencia tomarlo
	_____	_____	_____
	_____	_____	_____

Actividad física usa albuterol/levalbuterol ____ puffs, 15 minutos antes de la actividad con todas las actividades
 cuando siente que la necesita

Zona amarilla: Precaución

Síntomas: Algunos problemas de respiración – Tos, sibilancias, u opresión en el pecho – Problemas para trabajar o jugar – Se despierta en la noche

Medidor de flujo _____ a _____ (entre 50% y 79% de la mejor marca personal)

Medicamento/s de alivio rápido Albuterol/levalbuterol ____ puffs, cada 4 horas según sea necesario

Medicamento/s de control Continuar con los medicamentos de la Zona verde

Agregar _____ Cambiar por _____

El niño se debe sentir mejor dentro de 20–60 minutos del tratamiento de alivio rápido. Si el niño empeora o está en la Zona amarilla durante más de 24 horas, siga las instrucciones de la ZONA ROJA y llame al médico inmediatamente.

Zona roja: ¡Consigue ayuda ahora!

Síntomas: Muchos problemas de respiración – No puede trabajar o jugar – Empeora en vez de mejorar – El medicamento no ayuda **Medidor de flujo** _____ (menos del 50% de la mejor marca personal)

¡Tome el medicamento de alivio rápido AHORA! Albuterol/levalbuterol ____ puffs, _____
 (con qué frecuencia)

Llame al 911 inmediatamente si presenta las siguientes señales de peligro

- Dificultad para caminar/hablar debido a la falta de aire
- Labios o uñas azules
- Sigue en la zona roja después de 15 minutos

Contacto de emergencia Nombre _____ Teléfono (____) _____ - _____

Proveedor de atención médica Nombre _____ Teléfono (____) _____ - _____

Fecha ____ / ____ / ____

How to Use a Metered-Dose Inhaler with a Valved Holding Chamber (Spacer)

Prime a brand-new inhaler: Before using it for the first time, if you have not used it for more than 7 days, or if it has been dropped.



1. Shake inhaler 10 seconds.



2. Take the cap off the inhaler and valved holding chamber. Make sure the mouthpiece and valved holding chamber are clean and there is nothing inside the mouthpieces.



3. Put inhaler into the chamber/spacer.



4. Breathe out away from the device.



5. Put chamber mouthpiece in mouth.



6. Press inhaler once and breathe in deep and steadily.



7. Hold your breath for 10 seconds, then breathe out slowly.

If you need another puff of medicine, wait 1 minute and repeat steps 4-7.



8. Rinse with water and spit it out.

Proper inhalation technique is important when taking your asthma medicine(s) and monitoring your breathing. Make sure to bring all your medicines and devices to each visit with your primary care provider or pharmacist to check for correct use, or if you have trouble using them.

For more videos, handouts, tutorials and resources, visit [Lung.org](https://www.lung.org).

Scan the QR Code to access How-To Videos



You can also connect with a respiratory therapist for one-on-one, free support from the American Lung Association's Lung HelpLine at **1-800-LUNGUSA**.